

Physical Examples of Quantum Entropies: Properties, Calculations and Programmability

Stefanopoulos Dimitris

Aristotle's University of Thessaloniki
Subatomic Physics and Technological Applications

Thessaloniki, February 17, 2021

Table of contents

- 1 Intro
 - Motivation
 - Spectral theorem
 - Quantum Theory

- 2 Quantum entropies
 - von Neumann entropy
 - Other entropies
 - Quantum Relative entropy

- 3 Conclusions

Motivation

- Ill-defined conventions when calculating quantum entropies (for example $0 \ln 0 \equiv 0$) work operationally but it is not easy to use them when we automate the calculations with programming languages.
- It is not easy to find analytic calculations of quantum entropies in the literature.
- Why a physicist might care about quantum entropies?

Proposal for a Direct Measurement of the von Neumann Entropy and the Relative Entropy of Coherence

Bertúlio de Lima Bernardo^{1,2}

Variational approach to relative entropies (with application to QFT)

Stefan Hollands^{1*}

- Why do we even talk about more than one entropy? Different entropies can describe extra or different phenomena than the common definitions.

Quantum states and substates

Pure states are ket vectors $|\psi\rangle \in \mathcal{H}$. The density operator:

$$\rho = \sum_i p_i |\psi_i\rangle \langle \psi_i|$$

with $\text{Tr} \rho = 1, \rho = \rho^\dagger, \rho \geq 0$.

Note

The randomness of a quantum state expressed via the density matrix has two manifestations.

The reduced substate:

$$\rho^A = \text{Tr}_B (\rho^{AB}) = \sum_i (I_A \otimes \langle i|_B) \rho^{AB} (I_A \otimes |i\rangle_B)$$

$$\text{Tr}_B \left(\sum_{i,j,k,l} p_{ijkl} |i\rangle \langle k|_A \otimes |j\rangle \langle l|_B \right) = \sum_{i,j,k,l} p_{ijkl} |i\rangle \langle k|_A \text{Tr} (|j\rangle \langle l|_B)$$

Bell's inequalities

Based on EPR-like states $(|00\rangle + |11\rangle)/\sqrt{2}$ Bell discovered his famous inequalities.

ON THE EINSTEIN PODOLSKY ROSEN PARADOX

J. S. BELL[†]

Assuming a probability distribution of predictive properties \mathbb{P} consistent with *local realism* we are forced to find $E(\mathbb{P}) \leq 2$. However projective measurements lead to $E(\mathbb{P}) = 2\sqrt{2}$. Violation of these inequalities is though of as a sufficient criterion for entanglement.

Conditional entropies and their relation to entanglement criteria

Karl Gerd H. Vollbrecht[†] and Michael M. Wolf[†]

Entropic Bell Inequalities

N. J. Cerf[†] and C. Adami^{1,2}

Quantum information can be negative

Michal Horodecki¹, Jonathan Oppenheim² & Andreas Winter³

Negative entropy and information in quantum mechanics

N. J. Cerf[†] and C. Adami^{1,2}

Remark

- Negative conditional entropy is necessary condition for entanglement.
- Positive conditional entropy is a necessary condition for separability.

Basic theory and concepts

Definition

(Von Neumann entropy) The von Neumann entropy of a quantum state ρ is defined as:

$$S(\rho) \equiv -\mathbf{Tr}(\rho \log \rho).$$

Definition

(Heuristic von Neumann entropy) For a density matrix $\rho \in \mathcal{D}(\mathcal{H})$ the heuristic form of the von Neumann entropy is defined as:

$$S(\rho) = -\mathbf{Tr}(F(\rho))$$

in which F is the function $F : [0, 1] \rightarrow \mathbb{R}$:

$$F(x) = \lim_{\epsilon \rightarrow x} (\epsilon \log \epsilon)$$

For non-symbolic programming:

$$F(x) = \begin{cases} 0 & x = 0 \\ x \log x & x > 0 \end{cases}$$

Example-1

A statistical mixture of N orthogonal (mutually exclusive) pure states.

$$\rho = \sum_{i=0}^{N-1} \rho_i |i\rangle \langle i| = \text{diag}(\rho_0, \rho_1, \dots, \rho_{N-2}, \rho_{N-1})$$

$$S(\rho) = -\text{Tr} \left[\text{diag} \left(F(\rho_0), F(\rho_1), \dots, F(\rho_{N-2}), F(\rho_{N-1}) \right) \right] = - \sum_{i=0}^{N-1} \rho_i \log \rho_i$$

For a fixed temperature gas at a $T = 1/k_B\beta$ in a canonical ensemble model has a density operator:

$$\rho_{CE} = \frac{\exp(-\beta\hat{H})}{Z}$$

with β being a free parameter, \hat{H} and ϵ_n denotes the Hamiltonian and its eigenvalues and Z the quantum partition function:

$$Z = \text{Tr} \left(e^{-\beta\hat{H}} \right) = \sum_i e^{-\beta\epsilon_i}$$

based on the condition $\text{Tr}(\rho) = 1$. Thus:

$$\rho_i = e^{-\beta\epsilon_i} / Z$$

Conditional von Neumann entropy

Note

Definitions of classical probability theory can not be trivially used for non-commutative algebras.

Definition

(Conditional von Neumann Entropy) The quantum analog of the conditional entropy is defined as:

$$S(A | B) = S(AB) - S(B)$$

Where $S(AB) = S(\rho^{AB})$ and $S(B) = S(\text{Tr}_A \rho^{AB})$.

Example-II

Experimental investigation of partially entangled states for device-independent randomness generation and self-testing protocols

S. Gómez,^{1,2} A. Mattar,³ I. Machuca,^{1,2} E. S. Gómez,^{1,2} D. Cavalcanti,³ O. Jiménez Fariás,³ A. Acín,^{3,4} and G. Lima^{1,2}

¹*Departamento de Física, Universidad de Concepción, 160-C Concepción, Chile*

²*Millennium Institute for Research in Optics, Universidad de Concepción, 160-C Concepción, Chile.*

³*ICFO-Institut de Ciències Fotoniques, The Barcelona Institute*

of Science and Technology, 08860 Castelldefels, Barcelona, Spain

⁴*ICREA-Institució Catalana de Recerca i Estudis Avançats, Lluís Companys 23, 08010 Barcelona, Spain*

(Dated: February 5, 2019)

$$|\psi(\theta)\rangle = \cos\theta|00\rangle + \sin\theta|11\rangle, \quad 0 < \theta < \pi/2$$

Its density matrix:

$$\sigma^{AB}(\theta) = \begin{pmatrix} \cos^2\theta & 0 & 0 & \cos\theta\sin\theta \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \cos\theta\sin\theta & 0 & 0 & \sin^2\theta \end{pmatrix}$$

Example-II

$$\lambda_1 = 1, \lambda_2 = 0, \lambda_3 = 0, \lambda_4 = 0,$$

$$v_1 = \begin{pmatrix} \cot \theta \\ 0 \\ 0 \\ 1 \end{pmatrix}, v_2 = \begin{pmatrix} -\tan \theta \\ 0 \\ 0 \\ 1 \end{pmatrix}, v_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, v_4 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$$

Hence the modal matrix is:

$$M = \begin{pmatrix} \cot \theta & -\tan \theta & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix}$$

which gives:

$$\det(M) = -(\cos \theta \sin \theta)^{-1}.$$

As a result:

$$M^{-1} = \begin{pmatrix} \cos \theta \sin \theta & 0 & 0 & \sin^2 \theta \\ -\cos \theta \sin \theta & 0 & 0 & \cos^2 \theta \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

Example-II

So we have decomposed ρ as:

$$\sigma^{AB} = MDM^{-1}$$

in which $D = \text{diag}(\lambda_1, \lambda_2, \lambda_3, \lambda_4)$.

$$\begin{aligned} S(\sigma^{AB}) &= -\text{Tr}(F(\sigma^{AB})) \\ &= -\text{Tr}(F(MDM^{-1})) \\ &= -\text{Tr} \left[M \begin{pmatrix} F(1) & 0 & 0 & 0 \\ 0 & F(0) & 0 & 0 \\ 0 & 0 & F(0) & 0 \\ 0 & 0 & 0 & F(0) \end{pmatrix} M^{-1} \right] \\ &= 0. \end{aligned}$$

The result is expected since σ^{AB} is a pure state.

Remark

It can be generally proven that pure states have zero von Neumann entropy, using $\rho = \rho^2$.

Example-II

Let's trace out the second qubit using bra-ket notation and the linearity of the partial trace operator:

$$\begin{aligned}
 \sigma^A &= \text{Tr}_B(\sigma^{AB}) \\
 &= \text{Tr}_B \left(\cos^2 \theta |00\rangle \langle 00| + \cos \theta \sin \theta |00\rangle \langle 11| + \cos \theta \sin \theta |11\rangle \langle 00| + \sin^2 \theta |11\rangle \langle 11| \right) \\
 &= \text{Tr}_B \left(\cos^2 \theta |0\rangle \langle 0| \otimes |0\rangle \langle 0| + \cos \theta \sin \theta |0\rangle \langle 1| \otimes |0\rangle \langle 1| \right. \\
 &\quad \left. + \cos \theta \sin \theta |1\rangle \langle 0| \otimes |1\rangle \langle 0| + \sin^2 \theta |1\rangle \langle 1| \otimes |1\rangle \langle 1| \right) \\
 &= \cos^2 \theta |0\rangle \langle 0| \text{Tr}(|0\rangle \langle 0|) + \cos \theta \sin \theta |0\rangle \langle 1| \text{Tr}(|0\rangle \langle 1|) \\
 &\quad + \cos \theta \sin \theta |1\rangle \langle 0| \text{Tr}(|1\rangle \langle 0|) + \sin^2 \theta |1\rangle \langle 1| \text{Tr}(|1\rangle \langle 1|) \\
 &= \cos^2 \theta |0\rangle \langle 0| + \sin^2 \theta |1\rangle \langle 1| \\
 &= \begin{pmatrix} \cos^2 \theta & 0 \\ 0 & \sin^2 \theta \end{pmatrix}.
 \end{aligned}$$


Example-II

We see that the reduced density operator is diagonal. Hence, we don't need to decompose the matrix further. Let's calculate:

$$\begin{aligned}
 S(B | A)_\sigma &= S(\sigma^{AB}) - S(\sigma^A) \\
 &= 0 - S(\sigma^A) \\
 &= -S(\sigma^A) \\
 &= \mathbf{Tr}[F(\sigma^A)] \\
 &= \mathbf{Tr} \left[\begin{pmatrix} F(\cos^2 \theta) & 0 \\ 0 & F(\sin^2 \theta) \end{pmatrix} \right] \\
 &= 2 \sin^2 \theta \log(\sin \theta) + 2 \cos^2 \theta \log(\cos \theta)
 \end{aligned}$$

Example-II

This result demonstrates the general property of the quantum conditional entropy that pure entangled states have negative values. It is actually easy to see from the following plot that at the limits of $\theta \rightarrow 0$ and $\theta \rightarrow \pi/2$ the measure goes to 0.



We can easily see that the minimum is close to $-\log 2$. This is not an accident since it is common among the so called maximally entangled states. In particular, $\theta = \pi/4$ will give the maximal entanglement.

Example-III(Werner states)

Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model

Reinhard F. Werner*

Dublin Institute for Advanced Studies, 10 Burlington Road, Dublin 4, Ireland

(Received 1 May 1989)

Note on Separability of the Werner states in arbitrary dimensions

Arthur O. Pittenger †[‡] and Morton H. Rubin ‡

$$W^{[d^n]}(s) = (1 - s) \frac{1}{d^n} I + s |\Psi\rangle\langle\Psi|$$

where s is a free parameter, d is the dimension of the qudits, n is the number of qudits, $|\Psi\rangle$ is an entangled state and I is the identity matrix for the composed Hilbert space. It is proven that the state $W^{[d^n]}(s)$ is fully separable if and only if $s \leq (1 + d^{n-1})^{-1}$. We take the simplest case of $d = 2$, $n = 2$ and $|\Psi\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$ as prescribed.

Example-III(Werner states)


This state is separable iff $s \leq 1/3$:

$$W = \begin{pmatrix} (1+s)/4 & 0 & 0 & s/2 \\ 0 & (1-s)/4 & 0 & 0 \\ 0 & 0 & (1-s)/4 & 0 \\ s/2 & 0 & 0 & (1+s)/4 \end{pmatrix}$$

With

$$W^A = \begin{pmatrix} 1/2 & 0 \\ 0 & 1/2 \end{pmatrix}$$

$$\begin{aligned} S(B|A)_W &= S(AB) - S(A) \\ &= S(W) - S(W^A) \\ &= \frac{3}{4}(s-1) \log\left(\frac{1-s}{4}\right) - \frac{1}{4}(3s+1) \log\left(\frac{1}{4}(3s+1)\right) - \log(2) \end{aligned}$$



The blue point B has the coordinates $(1/3, 0.549306)$ and the red point $R(0.747614, 0)$ which was found via numerical methods. The diagram clearly illustrates that the state W while entangled ($s \geq 1/3$) has positive quantum conditional entropy, a fact emphasized by many sources. This particular example becomes negative for $s \gtrsim 0.747614$.

Renyi entropy

Definition

(Heuristic quantum Renyi entropy) For a density matrix $\rho \in \mathcal{D}(\mathcal{H})$ the heuristic form of the quantum Renyi entropy is defined as:

$$R(\alpha; \rho) = \frac{1}{1 - \alpha} \log \mathbf{Tr} (r(\alpha; \rho)), \alpha \in (0, 1) \cup (1, \infty)$$

in which r is the function $r : [0, 1] \rightarrow \mathbb{R}^+$: $r(\alpha; x) = x^\alpha$.

Remark

It is proven that: $\lim_{\alpha \rightarrow 1} R(\alpha; \rho) = S(\rho)$. For $\alpha \rightarrow 0$ and $\alpha \rightarrow \infty$ the Renyi entropy converges to the Hartley entropy and the Min entropy respectively.

Definition

(Quantum Conditional Renyi Entropy) The conditional form of the quantum Renyi entropy, for a density matrix $\rho \in \mathcal{D}(\mathcal{H})$ is defined as:

$$R(\alpha; A | B) = R(\alpha; \rho^{AB}) - R(\alpha; \rho^B)$$

Tsallis entropy

Definition

(Heuristic quantum Tsallis entropy) For a density matrix $\rho \in \mathcal{D}(\mathcal{H})$ the heuristic form of the quantum Tsallis entropy is defined as:

$$T(q; \rho) = \frac{1}{1-q} (\text{Tr} [t(q; \rho)] - 1), \quad q \in (0, 1) \cup (1, \infty)$$

in which t is the function $t : [0, 1] \rightarrow \mathbb{R}^+$: $t(q; x) = x^q$.

Remark

It can be proved that: $S(\rho) = \lim_{q \rightarrow 1} T(q; \rho)$. For $q \rightarrow 0$ and $q \rightarrow \infty$ the Tsallis entropy converges to $\text{rank}(\rho) - 1$ and 0 respectively.


Definition

(Quantum Conditional Tsallis Entropy) The conditional form of the quantum Tsallis entropy, for a density matrix $\rho \in \mathcal{D}(\mathcal{H})$ is defined as:

$$T(q; A | B) = \frac{T(q; \rho^{AB}) - T(q; \rho^B)}{1 + (1-q)T(q; \rho^B)}$$

Example-IV $(\rho(s) = s |0\rangle \langle 0| + (1 - s) |1\rangle \langle 1|)$


$$R(\alpha; \rho(s)) = \log [s^\alpha + (1 - s)^\alpha] / (1 - \alpha)$$



for $s = 0.5$ Renyi entropy takes the value $\log 2$ independent of α .

Example-IV $(\rho(s) = s |0\rangle \langle 0| + (1 - s) |1\rangle \langle 1|)$

$$T(q; \rho(s)) = [s^q + (1 - s)^q - 1] / (1 - q)$$



for $s = 0.5$ Tsallis entropy is not independent of q .

Example-V (Maximally mixed in N dimensions)

For

$$\rho = \sum_{i=0}^{N-1} \rho_i |i\rangle \langle i| = \text{diag}(\rho_0, \rho_1, \dots, \rho_{N-2}, \rho_{N-1}) = I_N/N$$

von Neumann, Renyi case:


$$S(\rho) = R(\alpha; \rho) = \log N \quad (1)$$

Tsallis case:

$$T(q; \rho) = \frac{1 - N^{1-q}}{q - 1} \quad (2)$$

Example-III(Werner states)


$$R(B|A)_W = \frac{\log(2^{1-\alpha}) - \log(4^{-\alpha}(3(1-s)^\alpha + (3s+1)^\alpha))}{\alpha - 1}$$



As an example, blue point B has the coordinates $(1, 0.978043)$ and the red point $R(1/3, 0.65241)$ which was found via numerical methods.

Example-III(Werner states)

$$T(B|A)_W = \frac{2^{-q-1} \left(-3(1-s)^q - (3s+1)^q + 2^{q+1} \right)}{q-1}$$



Blue point $B(0, 0.978043)$ and the red point $R(1/3, 0.826907)$.

Theory

Definition

(Quantum Relative Entropy) The quantum relative entropy $D(\rho\|\sigma)$ between density operators $\rho \in \mathcal{D}(\mathcal{H})$ and $\sigma \in \mathcal{L}(\mathcal{H})$ is defined by:

$$D(\rho\|\sigma) = \begin{cases} \text{Tr}[\rho \log \rho] - \text{Tr}[\rho \log \sigma] & \text{supp}(\rho) \subseteq \text{supp}(\sigma) \\ \infty & \text{otherwise} \end{cases}$$

Note

The quantum relative entropy is 0 iff $\rho = \sigma$.

Definition

(Heuristic Quantum Relative Entropy) The heuristic quantum relative entropy $Q(\rho\|\sigma)$ between the $d \times d$ density matrices ρ and σ is defined by:

$$Q(\rho\|\sigma) = S(\rho) - \lim_{\epsilon \rightarrow -\infty} \left(\text{Tr}[\rho G(\epsilon; \sigma)] \right)$$

in which $G : [0, 1] \rightarrow \mathbb{R}$ as:

$$G(\epsilon; x) = \begin{cases} \ln x & x \in (0, 1] \\ \epsilon & x = 0 \end{cases}$$

ϵ is a free parameter with $\epsilon \in \mathbb{R}$.

Conclusions

- It is tempting to conjecture that if there can be designed an experiment that measures $Q(\sigma||W)$ directly, we can detect and quantify the entanglement of the pure state $\sigma(\theta)$.
- It is proven for the quantum relative entropy that:

$$D(\rho||\sigma) = \lim_{\varepsilon \rightarrow 0^+} D(\rho||\sigma + \varepsilon I)$$

using a different function $G(\varepsilon; x) = x + \varepsilon$. Since different functions can be used to heuristically determine the quantum relative entropy, questions are raised regarding the class of functions $G(\varepsilon; x)$.

- Figures in slide 16,29 examples II,VI,VII do not exist in the current literature.